

1D Dynamics

1. MS p 38, Exercise 3. For more practice, find the fixed points of the following models. If there are no fixed points, say so. For each fixed point, analyse its stability using the derivative test. If the derivative test is inconclusive, say so.

- $x(t + 1) = Rx(t)$
- $x(t + 1) = a + x(t)$

2. Consider the model

$$\begin{aligned} X(t + 1) &= RX(t) \quad \text{if } X(t) < X_p \\ &= RX_p \exp(-b(X - X_p)) \quad \text{otherwise} \end{aligned}$$

- How many equilibria will there be if $R < 1$? If $R > 1$?
- Write down an equation that defines the equilibrium for the non-zero equilibrium if $R > 1$ and simplify it as far as you can (*don't try to solve it!*).
- For what values of the parameters is the zero equilibrium stable?
- Write down a criterion for the stability of the non-zero equilibrium. Substituting in from the question in step 2, can you express the stability criterion in terms of b and X^* ?

Linear models

3. We wish to model the volume of water that is present in a lake somewhere in Canada. Every year on New Year's Day, an engineer inspects the condition of the lake and makes an accurate calculation of the amount of water it contains. They started doing this on January 1st, 2014. At that time, the lake held exactly a million cubic meters of water. In the first year, they noticed the following three things:

- early February, when the snow melts, a quarter million cubic meters of water flows into the lake,

- due to evaporation and seepage during the warm summer months, the lake loses 20% of its water in July and 30% of its water in August,
- the September rains add half a million cubic meters of water to the lake.
- Let's call t the time in years, starting from $t = 0$ which corresponds to January 1st, 2014. Denote by $V(t)$ the amount of water (in millions of cubic meters) in the lake after t years. Remark that the question is already providing you with fact that $V(0) = 1$.

Now answer the following questions, in order.

- How much water will be present on January 1st, 2015? In other words, what is $V(1)$?
- What will be the amount $V(2)$ present on January 1st, 2016?
- Construct a recursion equation that predicts $V(t + 1)$ given $V(t)$.
- What kind of model is this? Classify it (univariate or multivariate, ...).
- Find the equilibrium amount of water in the lake.
- Characterize the stability of the equilibrium.

MLDD models

4. (Problem 3.5 in Mooney and Swift)

A copy machine is always in one of two states, either working or broken (not working). If it is working, there is a 70% chance that it will be working tomorrow. If it is broken there is a 50% change that it will still be broken tomorrow. Assume that one day is a natural time step.

- Draw a state diagram, labeling the states, using arrows to denote the direction of flow, and labeling paths with the per capita flow.
- Formulate the transition matrix.

- Assuming that the machine is working today, what is the probability that it will be working tomorrow? The next day?
- Explain what operations you would have to perform on \mathbf{M} to determine the long-term average probability that the machine is working on any given day.

5. (Problem 3.7 in Mooney and Swift). Do this problem, but don't do any of the actual linear algebra in part c: just identify the steps you would have to take (i.e. write down the numeric values of the matrices and identify what steps (inversion, multiplication, etc.) are necessary to solve the problem.

MNDD models

6. (From Jan Feys) Multivariate models also arise in economics. Consider the competition between two companies A and B in a **duopoly**, meaning that they have the whole market split between them. Company A brings x units to the market at a production cost of α per unit, while company B brings y at a production cost of β per unit. Assuming that the price the units sell for is $p = 1/(x + y)$, the profits (sales income minus production cost) of the companies are

$$P_A = \frac{x}{x + y} - \alpha x, \quad P_B = \frac{y}{x + y} - \beta y \quad .$$

Company A hired a mathematician to ensure maximum profitability. To this end, the mathematician sets out to maximize P_A by taking its derivative with respect to x and setting it to 0. He determines the optimal output to be

$$x = \sqrt{\frac{y}{\alpha}} - y \quad .$$

That means that company A decides how many units to produce purely on how many units company B is producing! Of course, company B also has a mathematician on payroll who comes to a similar conclusion. The discrete model that governs the dynamics of x and y is strongly coupled.

It reads

$$x(t+1) = \sqrt{\frac{y(t)}{\alpha}} - y(t),$$
$$y(t+1) = \sqrt{\frac{x(t)}{\alpha}} - x(t)$$

Assume α and β are positive constants. Answer the following three questions.

- Classify the model (according to the scheme we have been using in class: ULDD, etc.)
- What are the fixed points of the system?
- What is the stability of this system?

7. (Steve Walker) Find the equilibrium/equilibria of the following system and classify its/their stability.

$$x(t+1) = x(t) - y(t) + 1$$
$$y(t+1) = y(t) - y(t)^2 + x(t)$$

Python

8.

- Write a Python program that runs the discrete logistic map $X_{t+1} = RX_t(1 - X_t)$ for 100 time steps, i.e. from $t = 0$ up to $t = 99$, using parameter $R = 3$ and starting condition $X = 0.5$.
- Now write a new program that saves all the values from $X(0)$ to $X(99)$ in a numpy array.